

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
11 December 2003 (11.12.2003)

PCT

(10) International Publication Number  
**WO 03/101972 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 249/04**

(74) Agents: **NORTHROP, Thomas, E.** et al.; The Scripps Research Institute, 10550 North Torrey Pines Road, TPC-8, La Jolla, CA 92037 (US).

(21) International Application Number: PCT/US03/17311

(22) International Filing Date: 30 May 2003 (30.05.2003)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/385,041 30 May 2002 (30.05.2002) US

(71) Applicant (*for all designated States except US*): **THE SCRIPPS RESEARCH INSTITUTE** [US/US]; 10550 North Torrey Pines Road, La Jolla, CA 92037 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **SHARPLESS, Barry, K.** [US/US]; 7960 La Jota Way, La Jolla, CA 92037 (US). **FOKIN, Valery** [RU/US]; 935 Laguna Drive, #52, Carlsbad, CA 92008 (US). **ROSTOVSEV, Vsevolod** [RU/US]; 5 Kristina Court, Woodmill, Wilmington, DE 19808 (US). **GREEN, Luke** [GB/GB]; 94 Mawson Road, Cambridge, Cambridgeshire CB1 2EA (GB). **HIMO, Fahmi** [SE/SE]; Klippgatan 24A, S-171 47 Solna (SE).

**Published:**

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: COPPER-CATALYSED LIGATION OF AZIDES AND ACETYLENES

(57) Abstract: A metal catalyzed click chemistry ligation process is employed to bind azides and terminal acetylenes to give triazoles. In many instances, the reaction sequence regiospecifically ligates azides and terminal acetylenes to give only 1,4-disubstituted [1,2,3]-triazoles.



**WO 03/101972 A1**

## COPPER-CATALYSED LIGATION OF AZIDES AND ACETYLENES

### Description

#### Technical Field:

The invention relates to a stepwise Huisgen cycloaddition process catalyzed by copper(I). More particularly, the invention relates to a copper catalysed regioselective click chemistry ligation of azides and terminal alkynes to form triazoles.

#### Background:

Huisgen 1,3-dipolar cycloadditions are exergonic fusion processes that unite two unsaturated reactants (R. Huisgen, in *1,3-Dipolar Cycloaddition Chemistry*, (Ed.: A. Padwa), Wiley, New York, **1984**, pp. 1-176; and A. Padwa, in *Comprehensive Organic Synthesis*, (Ed.: B.M. Trost), Pergamon, Oxford, **1991**, Vol. 4, pp 1069-1109). For a review of asymmetric 1,3-dipolar cycloaddition reactions, see K.V. Gothelf, et al., *Chem. Rev.* **1998**, 98, 863-909. For a review of synthetic applications of 1,3-dipolar cycloadditions, see J. Mulzer, *Org. Synth. Highlights* **1991**, 77-95. Huisgen 1,3-dipolar cycloadditions provide fast access to an enormous variety of 5-membered heterocycles (a) W.-Q. Fan, et al., in *Comprehensive Heterocyclic Chemistry II*, (Eds.: A. R. Katritzky, et al.), Pergamon, Oxford, **1996**, Vol. 4, pp. 101-126; b) R.N. Butler, in *Comprehensive Heterocyclic Chemistry II*, (Eds.: A. R. Katritzky, et al.), Pergamon, Oxford, **1996**, Vol. 4, pp 621-678; and c) K. Banert, *Chem. Ber.* **1989**, 122, 911-918). The cycloaddition of azides and alkynes to give triazoles is arguably the most useful member of this family (a) R. Huisgen, *Pure Appl. Chem.* **1989**, 61, 613-628; b) R. Huisgen, et al., *Chem. Ber.* **1967**, 100, 2494-2507; c) W. Lwowski, in *1,3-Dipolar Cycloaddition Chemistry*, (Ed.: A. Padwa), Wiley, New York, **1984**; Vol. 1, Chapter 5; d) J. Bastide, et al., *Bull. Soc. Chim. Fr.* **1973**, 2555-2579; 2871-2887). However, probably because of concerns about the safety of working with organic azides, synthetic chemists, in both pure and applied fields, have not given this transformation the special attention it deserves. Although the actual cycloaddition

step may be faster and/or more regioselective for 1,3-dipoles other than azide, the latter is by far the most convenient to introduce and to carry hidden through many synthetic steps. Indeed, it appears to be the only three-atom dipole which is nearly devoid of side reactions.

5

Azides make only a fleeting appearances in organic synthesis, serving as one of the most reliable means to introduce a nitrogen substituent –  $R-X \rightarrow [R-N_3] \rightarrow R-NH_2$ . The azide intermediate is shown in brackets because it is generally reduced straightaway to the amine. Applications which leverage the unique reactivity offered by the azide group itself are disclosed by the following references from the laboratories of Aube, Banert, and Stoddart (a) P. Desai, et al., *J. Am. Chem. Soc.* **2000**, 122, 7226-7232; b) K. Banert, *Targets in Heterocyclic Systems* **1999**, 3, 1-32; K. Banert, *Liebigs Ann./Recl.* **1997**, 2005-18; c) J. Cao, et al., *J. Org. Chem.* **2000**, 65, 1937-46 and references cited therein. Although azide chemistry can be hazardous, the hazard of working with these reagents may be minimized by employing appropriate safety precautions. Azides are chemically important as a crucial functional group for click chemistry (H. C. Kolb, et al., *Angew. Chem. Int. Ed.* **2001**, 40, 2004-2021). The uniqueness of azides for click chemistry purposes arises from the extraordinary stability of these reagents toward  $H_2O$ ,  $O_2$ , and the majority of organic synthesis conditions. Indeed, organic azides, particularly in the aliphatic series, are exceptionally stable toward the common reactive chemicals, ranging from dioxygen and water to the aqueous solutions of highly-functionalized organic molecules which make up living cells. (E. Saxon, et al., *Science* **2000**, 287, 2007-2010; and K. L. Kiick, et al., *Proc. Natl. Acad. Sci. USA* **2002**, 99, 19-24). The spring-loaded nature of the azide group remains invisible unless a good dipolarophile is favorably presented. In fact, it was the razor sharp reactivity window for this cycloaddition process which spawned our “*in situ* click chemistry” ideas – an approach which resulted in discovery of the most potent non-covalent inhibitor of acetylcholinesterase known to date. (W. G. Lewis, et al., *Angew. Chem. Int. Ed.* **2002**, 41, 1053-1057). However, even then the desired triazole-forming cycloaddition may require elevated temperatures and, in any case, usually results in a mixture of the 1,4-

0

5

10

15

30

and 1,5- regioisomers (Fig 1A), unless the acetylene component is attached to an electron-withdrawing group such as a carbonyl or perfluoroalkyl (J. Bastide, et al., *Bull. Chim. Soc. Fr.* **1973**, 2294-2296; N. P. Stepanova, et al., *Zh. Org. Khim.* **1985**, 21, 979-983; N. P. Stepanova, et al., *Zh. Org. Khim.* **1989**, 25, 1613-1618; and D. Clarke, et al., *J. Chem. Soc. Perkin Trans. 1* **1997**, 1799-1804).

Efforts to control this 1,4- versus 1,5- regioselectivity problem have met with varying success (P. Zanirato, *J. Chem. Soc. Perkin Trans. 1* **1991**, 2789-2796; D. J. Hlasta, et al., *J. Org. Chem.* **1994**, 59, 6184-6189; C. A. Booth, et al., *Tet. Lett.* **1998**, 39, 6987-6990; S. J. Howell, et al., *Tetrahedron* **2001**, 57, 4945-4954; W. L. Mock, et al., *J. Org. Chem.*, **1989**, 54, 5302-5308; W. L. Mock *Top. Curr. Chem.* **1995**, 175, 1-24; J. Chen, et al., *Org. Lett.* **2002**, 4, 327-329; J. W. Wijnen, et al., *Tet. Lett.* **1995**, 36, 5389-5392; M. P. Repasky, et al., *Faraday Discuss.* **1998**, 110, 379-389).

In one report, copper (I) catalyzed regiospecific synthesis of peptidotriazoles was achieved in organic solvents using free azides and terminal acetylenes attached to a solid support. (C. W. Tornøe, et al., *J. Org. Chem.* **2002**, 67, 3057). Reactants were non-equimolar. An earlier report disclosed the formation, in the presence of copper (I), of a triazole, as a low yield byproduct, from a bifunctional reagent having an acetylene group and an in situ generated azide (G. L'abbe, *Bull. Soc. Chim. Belg.* 1984, 93, 579-592).

#### Summary:

One aspect of the invention is directed to a process for catalyzing a click chemistry ligation reaction between a first reactant having a terminal alkyne moiety and second reactant having an azide moiety for forming a product having a triazole moiety. More particularly, the click chemistry ligation reaction is catalyzed by an addition of Cu(II) in the presence of a reducing agent for reducing said Cu(II) to Cu(I), *in situ*, in catalytic amount. Preferred reducing agents include ascorbate, metallic copper, quinone, hydroquinone, vitamin K<sub>1</sub>, glutathione, cysteine, Fe<sup>2+</sup>, Co<sup>2+</sup>, and an applied electric potential. Further preferred reducing

agents include metals selected from the group consisting of Al, Be, Co, Cr, Fe, Mg, Mn, Ni, and Zn.

In an alternative aspect of the invention, a click chemistry ligation reaction between a first reactant having a terminal alkyne moiety and second reactant having an azide moiety for forming a product having a triazole moiety is catalyzed by performing the click chemistry ligation reaction in a solution in contact with metallic copper. The metallic copper contributes directly or indirectly to the catalysis of the click chemistry ligation reaction. In a preferred mode, the solution is an aqueous solution. The first and second reactants may be present during the click chemistry ligation reaction in equimolar amounts. Also, the click chemistry ligation reaction may be performed in a solution in contact, at least in part, with a copper vessel.

In another aspect of the invention, a click chemistry ligation reaction between a first reactant having a terminal alkyne moiety and second reactant having an azide moiety for forming a product having a triazole moiety is catalyzed by an addition of a catalytic amount of a metal salt having a metal ion selected from the group consisting of Au, Ag, Hg, Cd, Zr, Ru, Fe, Co, Pt, Pd, Ni, Rh, and W. In a preferred mode of this aspect of the invention, the click chemistry ligation reaction is performed in the presence of a reducing agent for reducing said metal ion to a catalytically active form. Preferred reducing agents include ascorbate, quinone, hydroquinone, vitamin K<sub>1</sub>, glutathione, cysteine, Fe<sup>2+</sup>, Co<sup>2+</sup>, an applied electric potential, and a metal selected from the group consisting of Al, Be, Co, Cr, Fe, Mg, Mn, Ni, and Zn.

In another aspect of the invention, a click chemistry ligation reaction between a first reactant having a terminal alkyne moiety and second reactant having an azide moiety for forming a product having a triazole moiety is performed in an aqueous solution and is catalyzed by a catalytic amount of copper (I). In a preferred mode of this aspect of the invention, the first and second reactants are present in equimolar amounts.

In another aspect of the invention, a click chemistry ligation reaction between a first reactant having a terminal alkyne moiety and second reactant having an azide moiety for forming a product having a triazole moiety is catalyzed by a catalytic amount of copper (I) while the first and second reactants are present in equimolar amounts. In a preferred mode of this aspect of the invention, the click chemistry ligation reaction is performed in an aqueous solution.


In another aspect of the invention, a click chemistry ligation reaction between a first reactant having a terminal alkyne moiety and second reactant having an azide moiety for forming a product having a triazole moiety is performed in a solution containing a catalytic amount of copper (I). However, in this instance, there is a proviso that either the first or second reactant is toxic or expensive and the remaining reactant is present in molar excess.

In another aspect of the invention, a click chemistry ligation reaction between a first reactant having a terminal alkyne moiety and second reactant having an azide moiety for forming a product having a triazole moiety is performed inside a cell. The cell is of a type that contains a catalytic quantity of copper(I) sufficient to catalyze the click chemistry ligation reaction. The copper(I) contributes to a catalysis of the click chemistry ligation reaction.

In another aspect of the invention, a click chemistry ligation reaction between a first reactant having a terminal alkyne moiety and second reactant having an azide moiety for forming a product having a triazole moiety is performed in a solvent containing a catalytic amount of a metal ion. The metal ions are selected from the group of metals consisting of Cu, Au, Ag, Hg, Cd, Zr, Ru, Fe, Co, Pt, Pd, Ni, Rh, and W. The metal ion contributes directly or indirectly to a catalysis of the click chemistry ligation reaction. The metal ion is coordinated to a ligand for solubilizing such metal ion within the solvent, for inhibiting oxidation of such metal ion, and for dissociating, in whole or in part, from such metal ion


during the catalysis of the click chemistry ligation reaction by said metal ion. A preferred ligand is acetonitrile. Another preferred ligand is a cyanide, nitrile, or isonitrile. Another preferred ligand is water. Other preferred ligands include nitrile, isonitrile, primary, secondary, or tertiary amine, a nitrogen bearing heterocycle, carboxylate, halide, alcohol, thiol, sulfide, phosphine, and phosphite. Other preferred ligands are polyvalent and include one or more functional groups selected from the group consisting of nitrile, isonitrile, primary, secondary, or tertiary amine, a nitrogen bearing heterocycle, carboxylate, halide, alcohol, thiol, sulfide, phosphine, and phosphite.

Another aspect of the invention is directed to a reactive intermediate for producing a product having triazole moiety. The reactive intermediate is represented by the following 6-membered ring structure:



In the above structure, R<sup>1</sup> is a 4-triazole substituent, R<sup>2</sup> is a 1-triazole substituent, L is a Cu ligand, and "n" is 1, 2, or 3.

Another aspect of the invention is directed to a reactive intermediate for producing a triazole. The reactive intermediate is represented by the following 6-membered ring structure:



In the above structure, R<sup>1</sup> is employable as a 4-triazole substituent, R<sup>2</sup> is employable as a 1-triazole substituent, L is a Cu ligand, and "n" is 1, 2, 3, or 4.

Another aspect of the invention is directed to a two step process of derivatizing an amine containing molecule with a triazole. In the first step of this

process the amine containing molecule is derivatized to form a terminal alkyne. Then, the product of the first step is ligated with an azide containing molecule by addition of the azide containing molecule in the presence of a catalytic amount of Cu to form a triazole derivative of the amine containing molecule.

Another aspect of the invention is directed to one step process for producing a polyvalent triazole. In this process, a polyazide core is derivatized by addition of a molecule having a terminal alkyne in the presence of a catalytic amount of Cu.

Another aspect of the invention is directed to a one step process for producing a polyvalent triazole. In this process, a polyalkyne core is derivatized by addition of an azide containing molecule in the presence of a catalytic amount of Cu.

#### Brief Description of Figures:

Figure 1A illustrates a prior art uncatalysed Huisgen cycloaddition of azides and terminal alkynes.

Figure 1B illustrates a copper catalysed regioselective ligation of azides and terminal alkynes.

Figure 2 illustrates proposed mechanism for the step-wise copper catalysed ligation of azides and terminal alkynes and compares this mechanism with a direct cycloaddition.

Figures 3A and 3B illustrate a table showing the synthesis of 1,4-disubstituted [1,2,3]-triazoles catalyzed by Cu<sup>I</sup> in the presence of ascorbate. All reactions were carried out in water with *t*-butanol as co-solvent, 0.25-0.5M in reactants, with 1 mol% of CuSO<sub>4</sub> and 10 mol% of sodium ascorbate, and were complete in 12-24 hours.



Figure 4 illustrates copper catalysed regioselective ligation of azides and terminal alkynes employing Cu(0) as a reducing agent to reduce Cu(2) to Cu(1).

Figure 5 illustrates use of human plasma as a solvent for the reaction. At 100 mM azide and 200 mM acetylene, the reaction is complete in 12-14 hours. At 2 mM azide and 4 mM acetylene, the reaction is approximately 80% complete after 48 hours. Note that the reaction does not decline in plasma, but merely slows down. Despite the high concentration of proteins and the binding of Cu to protein, the reaction proceeds well.

Figure 6 illustrates an exemplary two step procedure for derivatizing amine containing molecules, e.g. erythromycin, with triazoles. The procedure is applicable to any amine containing molecule.

Figure 7 illustrates a one step process using Cu catalysis for derivatizing polyazide cores to produce polyvalent triazoles.

Figure 8 illustrates a one step process using Cu catalysis for derivatizing polyalkyne cores to produce polyvalent triazoles.

#### Detailed Description:

The process is experimentally simple and appears to have enormous scope. While a number of copper(I) sources can be used directly (*vide infra*), it is disclosed that the catalyst is better prepared *in situ* by reduction of Cu<sup>II</sup> salts, which are less costly and often purer than Cu<sup>I</sup> salts (CuSO<sub>4</sub>•5H<sub>2</sub>O serves well). As the reductant, ascorbic acid and/or sodium ascorbate proved to be excellent, for they allow preparation of a broad spectrum of 1,4-triazole products in high yields and purity at 0.25 – 2 mol% catalyst loading. For a review of reactions of L-ascorbic acid with transition metals see M. B. Davies *Polyhedron* **1992**, *11*, 285-321 and references cited therein; redox properties of ascorbic acid are summarized in C. Creutz *Inorg. Chem.* **1981**, *20*, 4449. The reaction appears to

be very forgiving and does not require any special precautions. It proceeds to completion in 6 to 36 hours at ambient temperature in a variety of solvents, including aqueous *t*-butanol or ethanol and, very importantly, water with no organic co-solvent. Starting materials do not need to be dissolved in the reaction solvent. The reaction seems to proceed just as efficiently as long as adequate stirring is maintained. Although most experiments were performed at near neutral pH, the catalysis seems to proceed well at pH values ranging from ca. 4 to 12. The catalytic process is very robust and insensitive to usual reaction parameters.

It is further disclosed that Cu<sup>0</sup> can also be used as a source of the catalytic species. Although these reactions may take longer to proceed to completion, the experimental procedure is exceedingly simple. For example, bis-triazole shown in entry 2 (Figure 3A) was obtained in quantitative yield after stirring the corresponding azide and acetylene components for 24 h with ca. 2 g of copper metal turnings. The turnings were removed at the end of the reaction, and the pure white product was collected by simple filtration.

The reaction between phenyl propargyl ether and benzylazide in the presence of 5 mol% of sodium ascorbate and 1 mol% of copper(II) sulfate in 2:1 water/*t*-butanol mixture furnished the 1,4-disubstituted triazole product in 91% yield after stirring for 8 hours at room temperature in a capped scintillation vial, but otherwise with no effort to exclude oxygen [eq. (2)]. The regiochemistry of the product was established by NOE experiments and confirmed by the X-ray crystallographic analysis.

For comparison, the thermal reaction (neat, 92 °C, 18h) between these substrates gives both regioisomers in a ratio of 1.6:1 in favor of the 1,4-isomer, as illustrated in Figure 1A.

The scope of this copper-catalyzed triazole synthesis is partly revealed by the examples in Figures 3A and B; note especially the lack of functional group

interference. These triazoles are obtained using a procedure which generally involves little more than stirring the reagents and filtering off pure products. Various substituted primary, secondary, tertiary, and aromatic azides readily participate in this transformation. Tolerance for variations in the acetylene component is also excellent.

Copper(I) salts, for example CuI, CuOTf•C<sub>6</sub>H<sub>6</sub> and [Cu(NCCH<sub>3</sub>)<sub>4</sub>]PF<sub>6</sub>, can also be used directly in the absence of a reducing agent. These reactions usually require acetonitrile as co-solvent and one equivalent of a nitrogen base (e.g. 2,6-lutidine, triethylamine, diisopropylethylamine, or pyridine). However, formation of undesired byproducts, primarily diacetylenes, bis-triazoles, and 5-hydroxytriazoles, was often observed. For a recent summary of the reactions of copper(I) complexes with dioxygen, see S. Schindler *Eur. J. Inorg. Chem.* **2000**, 2311-2326 and A. G. Blackman, W. B. Tolman in *Structure and Bonding*, B. Meunier, Ed., Springer-Verlag, Berlin, Heidelberg, **2000**, vol. 97, p. 179-211. This complication with direct use of Cu<sup>I</sup>-species was minimized when 2,6-lutidine was used, and exclusion of oxygen further improved product purity and yield. For example, ethylpropiolate and benzyl azide furnished the corresponding 1,4-triazole in 55% yield when this procedure was used, but only trace amount of the product was obtained with 1 equiv. of triethylamine and no exclusion of oxygen. Even though a broad range of both acetylene and azide components react readily in the acetonitrile system, we prefer the even more reliable and simple Cu<sup>II</sup>/ascorbate aqueous system (with or without co-solvents and amine buffers/additives).

A mechanistic proposal for the catalytic cycle is illustrated in Figure 2. It begins unexceptionally with formation of the copper(I) acetylide **i** (G. van Koten, J. G. Noltes in *Comprehensive Organometallic Chemistry*, G. Wilkinson, Ed., vol. 2, chap. 14, Pergamon Press, **1982**, 720). As expected, no reaction is observed with internal alkynes. It is disclosed herein that extensive density functional theory calculations offer compelling evidence which strongly disfavors, by approximately 12-15 kcal, the concerted [2+3] cycloaddition (B-direct) and points to a stepwise, annealing sequence (B1 → B2 → B3), which proceeds via the 6-membered copper

containing intermediate **iii** (M. P. Doyle, et al., in *Modern Catalytic Methods for Organic Synthesis with Diazo Compounds* Wiley (New York), **1997**, 163-248). Hence, the term 'ligation' is employed herein to denote the step-wise [2+3] cycloaddition catalyzed by copper(I).

5

The Cu<sup>I</sup>-catalyzed transformation described here – a high-yielding and simple to perform 'fusion' process leading to a thermally and hydrolytically stable triazole connection – is an ideal addition to the family of click reactions. The process exhibits broad scope and provides 1,4-disubstituted [1,2,3]-triazole products in excellent yields and near perfect regioselectivity. The reactivity of copper(I) acetylides with organic azides is disclosed herein to be effectively unstoppable.

0

This new catalytic process offers an unprecedented level of selectivity, reliability and scope for those organic synthesis endeavors which depend on the creation of covalent links between diverse building blocks. Several applied projects which highlight the capabilities of the process are illustrated in Figures 6-8.

15


#### Experimental procedure:

General procedure as exemplified for the synthesis of 17-[1-(2,3-dihydroxypropyl)-1H-[1,2,3]triazol-4-yl]-estradiol. 17-ethynyl estradiol (888 mg, 3 mmol) and (S)-3-azidopropane-1,2-diol (352 mg, 3 mmol) were suspended in 12 mL of 1:1 water/*t*-butanol mixture. Sodium ascorbate (0.3 mmol, 300  $\mu$ L of freshly prepared 1M solution in water) was added, followed by copper(II) sulfate pentahydrate (7.5 mg, 0.03 mmol, in 100  $\mu$ L of water). The heterogeneous mixture was stirred vigorously overnight, at which point it cleared and TLC analysis indicated complete consumption of the reactants. The reaction mixture was diluted with 50 mL of water, cooled in ice, and the white precipitate was collected by filtration. After washing with cold water (2X25 mL), the precipitate was dried under vacuum to afford 1.17 g (94%) of pure product as off-white powder. M.p. 228–230 °C. Elemental analysis calculated: C 64.02%, H 7.71%, N 9.74%; found: C 64.06%, H 7.36%, N 9.64%.

20

25

30



### Reducing environment effect:

0 Cu(I) is very easily oxidized to Cu(II) – primarily by oxygen, but even milder oxidants can accomplish this. Additionally, Cu(I) mediates oxidative coupling of acetylenes and other organocopper species, which leads to reduced yields and contaminated products. All these problems can be circumvented by the use of a mild reducing agent. Various reducing agents can be used: ascorbate,

5 hydroquinone, other quinones (such as vitamin K<sub>1</sub>), copper turnings/wire, glutathione, cysteine, Fe<sup>2+</sup>, Co<sup>2+</sup>, etc. Virtually any reductant may be employed that is not so powerful so as to rapidly reduce Cu(II) to Cu(0).

### Ligands: Acetonitrile effect:


0 Metals do not exist in solutions “naked” – there are always ligands present in the coordination sphere, be it solvent molecules or ‘designer’ ligands. Ligands have a profound effect on reactivity of the metal center, as well as red/ox properties of the metal: (a) they can stabilize a certain oxidation state of the metal, e.g. Cu(I) is the desirable form in our case, and (b) just as importantly, they can

15 keep the catalytic species in the solution, thereby making it constantly available for the catalysis. Both of these requirements have to be fulfilled in order for a metal/ligand combination to be useful in catalyzing a desired transformation cycloaddition, or ligation, in the present instance.


20 Copper-mediated organic transformations have been studied for over 70 years, and the literature on the subject is quite extensive. An important lesson of the prior art is that cyanides and/or nitriles are some of the best ligands for Cu(I),

which is usually tetracoordinated, forming tetrahedral complexes. In fact, acetonitrile coordinates to Cu(I) so strongly that  $[\text{Cu}(\text{CH}_3\text{CN})_4]^+\text{PF}_6^-$  complex is a commercially available, oxygen-insensitive Cu(I) preparation (i.e., this Cu(I) is *unreactive*). This 'overstabilization' is clearly a liability when *reactivity* is our goal.

The reaction indicated below illustrates the point. When water/alcohol mixtures are employed as solvents (note that both are weak ligands for Cu(I)), the reaction is complete in under 6 hrs. However, when acetonitrile is used as a solvent, no reaction is observed at all even after 24 hrs under otherwise identical conditions.



To explain this phenomenon, recall the mechanism of the reaction. In order for the ligation to proceed, the azide must coordinate to the copper (step B1) and one ligand must be removed from the coordination sphere of the metal, and in case of a strongly coordinated acetonitrile, this step is disfavored.



Therefore, in order to have a useful reactivity window, one should choose or design such ligands that do bind to the metal relatively well, keep it in the correct oxidation state and in solution (i.e. not aggregated to the point of forming a polymeric precipitate), but in the same time can come off the metal center to allow formation of intermediate **ii**, which is a necessary step in the catalytic sequence. To use the example in hand, addition of an amine, such as triethylamine or 2,6-lutidine to the acetonitrile system described above, solves the problem of reactivity – the product is formed in quantitative yield after ca. 8 hrs.

Preferred ligands include cyanides, nitriles, isonitriles, primary or secondary amines, carboxylates, halides, alcohols, and thiols. Chloride is the preferred halide and best employed at 1-5 M. Polyvalent ligands that include one or more functional groups selected from nitrile, isonitrile, primary or secondary amine, carboxylate, alcohol, and thiol may also be employed.

#### Other Metal Catalysts:

Cu is not the only metal capable of catalyzing this type of cycloaddition. As long as intermediate **ii** can be formed (i.e. the metal is or can become coordinatively unsaturated), other metals known to form stable acetylides may also be employed. Exemplary metals that can form stable acetylides include Cu, Au, Ag, Hg, Cd, Zr, Ru, Fe, Co, Pt, Pd, Ni, Rh, and W. It is a matter of finding the right metal/ligand combination. Copper is unique among other metals because it is so well supported in aqueous systems, which results in exceedingly simple experimental procedure and amazing functional group tolerance.

#### Catalysis of ligation reaction by Metallic Container:

Metallic containers can also be used as a source of the catalytic species to catalyze the ligation reaction. For example, a copper container ( $\text{Cu}^0$ ) may be employed to catalyze the reaction. In order to supply the necessary ions, the reaction solution must make physical contact with the a copper surface of the

container. Alternatively, the reaction may be run in a non-metallic container, and the catalytic metal ions supplied by contacting the reaction solution with a copper wire, copper shavings, or other structures. Although these reactions may take longer to proceed to completion, the experimental procedure is exceedingly simple. For example, bis-triazole shown in entry 2 (Figure 3A) was obtained in quantitative yield after stirring the corresponding azide and acetylene components for 24 h with ca. 2 g of copper metal turnings. The turnings were removed at the end of the reaction, and the pure white product was collected by simple filtration.

#### Alternative Reducing Agents:

Metals may be employed as reducing agents to maintain the oxidation state of the Cu (I) catalyst or of other metal catalysts. Preferred metallic reducing agents include Cu, Al, Be, Co, Cr, Fe, Mg, Mn, Ni, and Zn. Alternatively, an applied electric potential may be employed to maintain the oxidation state of the catalyst.

#### In vivo Catalysis:

The reaction proceeded well in fresh human plasma (protein loading 65 – 85 mg/mL,  $C_{\text{azide}} = C_{\text{alkyne}} = 5\text{mM}$ ;  $C_{\text{Cu}^+} = 100\text{ mM}$ ) and in whole blood, indicating that copper species remained available for the catalysis despite being heavily bound to plasma proteins and indicating that the ligation reaction can be catalyzed by copper and/or other metals ions and templates *in vivo*, including intracellularly. The reaction proceeds fresh human plasma and intracellularly in whole blood without noticeable loss of catalytic turnover and without noticeable protein precipitation.



Cu(I) salt used directly:

If Cu(I) salt is used directly, no reducing agent is necessary, but acetonitrile or one of the other ligands indicate above should be used as a solvent (to prevent rapid oxidation of Cu(I) to Cu(II) and one equivalent of an amine should be added (to accelerate the otherwise extremely slow reaction – *vide supra*). In this case, for better yields and product purity, oxygen should be excluded. Therefore, the ascorbate (or any other reducing) procedure is often preferred over the unreduced procedure. The use of a reducing agent is procedurally simple, and furnishes triazole products in excellent yields and of high purity.

What is claimed is:

1. A process comprising the following step:  
catalyzing a click chemistry ligation reaction between a first reactant having a terminal alkyne moiety and second reactant having an azide moiety for forming a product having a triazole moiety, the click chemistry ligation reaction being catalyzed by an addition of Cu(II) in the presence of a reducing agent for reducing said Cu(II) to Cu(I), *in situ*, in catalytic amount.
2. A process according to claim 1 wherein the reducing agent is selected from the group consisting of ascorbate, quinone, hydroquinone, vitamin K<sub>1</sub>, metallic copper, glutathione, cysteine, Fe<sup>2+</sup>, Co<sup>2+</sup>, and an applied electric potential.
3. A process according to claim 1 wherein the reducing agent is a metal selected from the group consisting of Cu, Al, Be, Co, Cr, Fe, Mg, Mn, Ni, and Zn.
4. A process comprising the following step:  
catalyzing a click chemistry ligation reaction between a first reactant having a terminal alkyne moiety and second reactant having an azide moiety for forming a product having a triazole moiety, the click chemistry ligation reaction being performed in a solution in contact with metallic copper, said metallic copper contributing directly or indirectly to the catalysis of the click chemistry ligation reaction.
5. A process according to claim 4 wherein the solution is an aqueous solution.
6. A process according to claim 4 wherein the first and second reactants are present during the click chemistry ligation reaction in equimolar amounts.
7. A process according to claim 4 wherein the click chemistry ligation reaction is performed in a solution in contact, at least in part, with a copper vessel.

8. A process comprising the following step:  
catalyzing a click chemistry ligation reaction between a first reactant having a terminal alkyne moiety and second reactant having an azide moiety for forming a product having a triazole moiety, the click chemistry ligation reaction being catalyzed by an addition of a catalytic amount of a metal salt having a metal ion selected from the group consisting of Au, Ag, Hg, Cd, Zr, Ru, Fe, Co, Pt, Pd, Ni, Rh, and W.
9. A process according to claim 8 wherein the click chemistry ligation reaction being performed in the presence of a reducing agent for reducing said metal ion to a catalytically active form.
10. A process according to claim 9 wherein the reducing agent is selected from the group consisting of ascorbate, quinone, hydroquinone, vitamin K<sub>1</sub>, glutathione, cysteine, Fe<sup>2+</sup>, Co<sup>2+</sup>, an applied electric potential, and a metal, the metal being selected from the group consisting of Al, Be, Co, Cr, Fe, Mg, Mn, Ni, and Zn.
11. A process comprising the following step:  
catalyzing a click chemistry ligation reaction between a first reactant having a terminal alkyne moiety and second reactant having an azide moiety for forming a product having a triazole moiety, the click chemistry ligation reaction being performed in an aqueous solution and being catalyzed by a catalytic amount of copper (I).
12. A process according to claim 11 wherein the first and second reactants are present in equimolar amounts.
13. A process comprising the following step:  
catalyzing a click chemistry ligation reaction between a first reactant having a terminal alkyne moiety and second reactant having an azide moiety for forming a product having a triazole moiety, the click chemistry ligation

reaction being catalyzed by a catalytic amount of copper (I), said first and second reactants being present in equimolar amounts.

14. A process according to claim 13 wherein the click chemistry ligation reaction is performed in an aqueous solution.

15. A process comprising the following step:

catalyzing a click chemistry ligation reaction between a first reactant having a terminal alkyne moiety and second reactant having an azide moiety for forming a product having a triazole moiety, the click chemistry ligation reaction being performed in a solution containing a catalytic amount of copper (I);

with a proviso that either the first or second reactant is toxic or expensive and the remaining reactant is present in molar excess.

16. A process comprising the following step:


catalyzing a click chemistry ligation reaction between a first reactant having a terminal alkyne moiety and second reactant having an azide moiety for forming a product having a triazole moiety, the click chemistry ligation reaction being performed inside a cell, said cell containing a catalytic quantity of copper(I) sufficient to catalyze the click chemistry ligation reaction, said copper(I) contributing to a catalysis of the click chemistry ligation reaction.

17. A process comprising the following step:

catalyzing a click chemistry ligation reaction between a first reactant having a terminal alkyne moiety and second reactant having an azide moiety for forming a product having a triazole moiety, the click chemistry ligation reaction being performed in a solvent containing a catalytic amount of a metal ion selected from ions from the group of metals consisting of Cu, Au, Ag, Hg, Cd, Zr, Ru, Fe, Co, Pt, Pd, Ni, Rh, and W, said metal ion contributing directly or indirectly to a catalysis of the click chemistry ligation

reaction, said metal ion being coordinated to a ligand for solubilizing said metal ion within said solvent, for inhibiting oxidation of said metal ion, and for dissociating, in whole or in part, from said metal ion during the catalysis of the click chemistry ligation reaction by said metal ion.


18. A process according to Claim 17 wherein said ligand is acetonitrile.
19. A process according to Claim 17 wherein said ligand is a cyanide, nitrile, or isonitrile.
20. A process according to Claim 17 wherein said ligand is water.
21. A process according to Claim 17 wherein said ligand is selected from the group consisting of nitrile, isonitrile, primary, secondary, or tertiary amine, a nitrogen bearing heterocycle, carboxylate, halide, alcohol, thiol, sulfide, phosphine, and phosphite.
22. A process according to Claim 17 wherein said ligand is polyvalent and includes one or more functional groups selected from the group consisting of nitrile, isonitrile, primary, secondary, or tertiary amine, a nitrogen bearing heterocycle, carboxylate, halide, alcohol, thiol, sulfide, phosphine, and phosphite.
23. A reactive intermediate for producing a product having triazole moiety, the reactive intermediate being represented by the following 6-membered ring structure:



wherein:

- R<sup>1</sup> is a 4-triazole substituent,
- R<sup>2</sup> is a 1-triazole substituent,
- L is a Cu ligand, and
- "n" is 1, 2, or 3.

24. A reactive intermediate for producing a triazole, the reactive intermediate being represented by the following structure:



wherein:

$\text{R}^1$  is a 4-triazole substituent,

$\text{R}^2$  is a 1-triazole substituent,

L is a Cu ligand, and

"n" is 1, 2, 3, or 4.

25. A two step process of derivatizing an amine containing molecule with a triazole, the process comprising the following steps:

Step A: derivatizing the amine containing molecule to form a terminal alkyne, then

Step B: ligating the product of said Step A with an azide containing molecule by addition of the azide containing molecule in the presence of a catalytic amount of Cu to form a triazole derivative of the amine containing molecule.

26. A one step process for producing a polyvalent triazole, the process comprising the following step:

derivatizing a polyazide core by addition of a molecule having a terminal alkyne in the presence of a catalytic amount of Cu.

27. A one step process for producing a polyvalent triazole, the process comprising the following step:

derivatizing a polyalkyne core by addition of an azide containing molecule in the presence of a catalytic amount of Cu.




Figure 1A





Figure 1B





**Figure 3A**






Entry	Product	yield, %
1		94
2		93
3		82
4		84
5		92

Figure 3B







Entry	Product	yield, %
6		92
7		84
8		88
9		90
10		80
11		94

Figure 4






Figure 5




Human plasma as a solvent

Figure 6




Facile derivatization of complex molecules:  
Erythromycin

Figure 7



Polyvalency: exemplary polyazide cores

Figure 8



Polyvalency: exemplary alkyne cores

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/17311

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>																										
IPC(7) : C07D 249/04 US CL : 548/255																										
According to International Patent Classification (IPC) or to both national classification and IPC																										
<b>B. FIELDS SEARCHED</b>																										
Minimum documentation searched (classification system followed by classification symbols) U.S. : 548/255																										
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched																										
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Please See Continuation Sheet																										
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>																										
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.																								
A	LEWIS ET AL. Click Chemistry In Situ: Acetylcholinesterase as a Reaction Vessel for the Selective Assembly of a Femtomolar Inhibitor from an Array of Building Blocks. Angewandte Chemie International Edition, 2002, Vol. 41, No. 6, pages 1053-1057.	2																								
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.																										
<table border="0"> <tr> <td colspan="2">* Special categories of cited documents:</td> <td>"T"</td> <td>later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>"A"</td> <td>document defining the general state of the art which is not considered to be of particular relevance</td> <td>"X"</td> <td>document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>"E"</td> <td>earlier application or patent published on or after the international filing date</td> <td>"Y"</td> <td>document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>"L"</td> <td>document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>"&amp;"</td> <td>document member of the same patent family</td> </tr> <tr> <td>"O"</td> <td>document referring to an oral disclosure, use, exhibition or other means</td> <td></td> <td></td> </tr> <tr> <td>"P"</td> <td>document published prior to the international filing date but later than the priority date claimed</td> <td></td> <td></td> </tr> </table>			* Special categories of cited documents:		"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	"A"	document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	"E"	earlier application or patent published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&"	document member of the same patent family	"O"	document referring to an oral disclosure, use, exhibition or other means			"P"	document published prior to the international filing date but later than the priority date claimed		
* Special categories of cited documents:		"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention																							
"A"	document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone																							
"E"	earlier application or patent published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art																							
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&"	document member of the same patent family																							
"O"	document referring to an oral disclosure, use, exhibition or other means																									
"P"	document published prior to the international filing date but later than the priority date claimed																									
Date of the actual completion of the international search 29 July 2003 (29.07.2003)		Date of mailing of the international search report <b>21 AUG 2003</b>																								
Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US Commissioner for Patents P.O. Box 1450 Alexandria, Virginia 22313-1450 Facsimile No. (703)305-3230		Authorized officer <i>Rebecca L. Anderson</i> Telephone No. (703) 308-1235																								



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/17311

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claim Nos.: 1 and 3-27  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
Please See Continuation Sheet
3. ☐ Claim Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
  2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
  3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
- Remark on Protest ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

PCT/US03/17311

## Continuation of Box I Reason 2:

In these claims, the numerous variables (e.g. the various possible substituents on the terminal alkyne, on the azide and on the triazole and the various possible reducing agents, etc.) and their voluminous complex meanings and their seemingly endless permutations and combinations make it virtually impossible to determine the full scope and complete meaning of the claimed subject matter. As presented, the claimed subject matter cannot be regarded as being a clear and concise description for which protection is sought and as such the listed claims do not comply with the requirements of PCT article 6. Thus it is impossible to carry out a meaningful search on same. A search will be made on the first discernable invention of claim 2 which is the experimental procedure as found on pages 11 and 12 of the instant specification.

## Continuation of B. FIELDS SEARCHED Item 3:

CAS ONLINE

CASREACT reaction search